arXiv:1403.6798v1 [math.GR] 26 Mar 2014

Splitting automorphisms of prime power orders of free Burnside groups

V. S. Atabekyan

March 27, 2014

Abstract

We prove that if the order of a splitting automorphism of free Burnside group $B(m,n)$ of odd period $n \geq 1003$ is a prime power, then the automorphism is inner. Thus, we give an affirmative answer to the question on the coincidence of splitting and inner automorphisms of free Burnside groups $B(m,n)$ for automorphisms of orders p^k (p is a prime number). This question was posed in the Kourovka Notebook in 1990 (see 11th ed., Question 11.36. b).

Keywords: Splitting automorphism, inner automorphism, normal automorphism, free Burnside group

MSC: 20B27, 20F50, 20E36, 20F28

1 Introduction

An automorphism φ of a group G is called *a splitting automorphism of period n*, if $\varphi^n = 1$ and $g\, g^{\varphi} g^{\varphi^2} \cdots g^{\varphi^{n-1}} = 1$ for all $g \in G$. Various authors studied groups with splitting automorphism. The known theorem of O.Kegel, proved in 1961, states that any finite group possessing a nontrivial splitting automorphism of prime order is nilpotent (see [8]). E.Khukhro proved that any solvable group possessing a nontrivial splitting automorphism of prime order is also nilpotent (see [9]). E.Jabara in [10] established that a finite group with the splitting automorphism of order 4 is solvable.

It is not difficult to check that if φ is a splitting automorphism of period n of G, then $g^{\varphi^{n-1}} \cdots g^{\varphi^2} g^{\varphi}\, g = 1$ for any $g \in G$. Let φ be a splitting automorphism of period n of some group G. Since $(\varphi\, g)^n = \varphi^n g^{\varphi^{n-1}} \cdots g^{\varphi^2} g^{\varphi}\, g$, this precisely means that the relation $(\varphi\, g)^n = 1$

[0]The author was supported in part by State Committee Science MES RA grant in frame of project 13-1A246

holds in the holomorph $Hol(G)$ of G for each $g \in G$. In particular, the identity automorphism of a given group G is a splitting automorphism of period n if and only if the identity $x^n = 1$ holds in G.

It is easy to check that if in a given group G the identity $x^n = 1$ is valid, then each inner automorphism of G is an inner splitting automorphism of period n. However, the converse statement is false. Indeed, let F_2 be an absolutely free group with free generators a and b and let C_n be a cyclic group of order n with generator element c. Let us consider the group $\Gamma = F_2/F_2^n \times C_n$ of period n, where F_2^n stands for the subgroup generated by all possible nth powers of the elements of F_2. It can readily be seen that for any $n > 1$, the automorphism $\alpha : \Gamma \to \Gamma$ given on the generators by the formulae $\alpha(a) = ac$, $\alpha(b) = bc$, and $\alpha(c) = c$, is a splitting automorphism of period n. However, this is an outer automorphism because it is clear that no relation of the form $u^{-1}au = ac$ can hold in Γ.

S.V.Ivanov in [11] posed the following problem: *Let n be large enough odd number and $m > 1$. Is it true that each automorphism φ of $B(m, n)$, that satisfies the relations $\varphi^n = 1$ and $g\, g^{\varphi} g^{\varphi^2} \cdots g^{\varphi^{n-1}} = 1$ for any g in $B(m, n)$, is inner* (see [11], Question 11.36. b))? In fact, Ivanov's problem concerns splitting automorphisms of period n of the groups $B(m, n)$. By definition, the free Burnside group $B(m, n)$ of period n and rank m has the following presentation

$$B(m, n) = \langle a_1, a_2, ..., a_m \mid X^n = 1 \rangle,$$

where X ranges over the set of all words in the alphabet $\{a_1^{\pm 1}, a_2^{\pm 1}, \ldots, a_m^{\pm 1}\}$. The group $B(m, n)$ is a quotient group of the free group F_m of rank m by normal subgroup F_m^n, generated by all nth powers of the elements of F_m. Every periodic group of period n with m generators is a quotient group of $B(m, n)$.

The main Theorem 2.1 of the present paper gives a positive answer to the above mentioned question for all splitting automorphisms of prime power order of free Burnside groups $B(m, n)$ for odd periods $n \geq 1003$. In the paper [6], we established some properties of splitting automorphisms of the groups $B(m, n)$. Those properties have proved useful in our proof of the main result 2.1. Theorem 2.1 strengthens the result of [6], where the Ivanov's problem was solved only for automorphisms of prime orders.

1.1 On simple periodic groups

S.I.Adian and I.G.Lysenok in [2] proved, that for any $m > 1$ and odd $n \geq 1003$ there exists a maximal normal subgroup N of free Burnside group $B(m, n)$ such that the quotient group $B(m, n)/N$ is an infinite group, every proper subgroup of which is contained in some cyclic subgroup of order n. The groups constructed in [2] are infinite simple groups in which the

identity relation $x^n = 1$ holds. This groups are called 'Tarski monsters' since Tarski has formulated a question on the existence of such groups. The first examples of Tarski monsters were constructed by A.Yu.Olshanskii [12] for prime periods $n > 10^{75}$. It is now known that for every odd $n \geq 1003$ there are continuum many non-isomorphic Tarski-monsters of period n (see [3], [13, Theorem 28.7], [4]). Each of these Tarski monsters is a result of a factorization of the group $B(m, n)$ by some maximal normal subgroup of, which leads to an infinite group containing only cyclic proper subgroups.

We denote by $\mathcal{M}_n$ the set of all normal subgroups $N \lhd B(m, n)$ for which the quotient group $B(m, n)/N$ is a Tarski monster.

The following two statements were proved in [5] (see also [7]) and [6] respectively. They play the key role in the proof of main Theorem 2.1.

LEMMA **1.1.** (see [5, Corrolary 2]) *Let $n \geq 1003$ be an odd number and let φ be an automorphism of $B(m, n)$ such that for any normal subgroup $N \in \mathcal{M}_n$ the equality $N^\varphi = N$ holds. Then φ is an inner automorphism.*

LEMMA **1.2.** (see [6, Lemma 4]) *If φ is an arbitrary nontrivial splitting automorphism of period n of the group $B(m, n)$, where $n \geq 1003$ is odd, then the stabilizer of any normal subgroup $N \in \mathcal{M}_n$ under the action of the cyclic group $\langle\varphi\rangle$ is nontrivial.*

2 The main result

THEOREM **2.1.** *Let ϕ be a splitting automorphism of period n of the group $B(m, n)$, where $n \geq 1003$ is an odd number. If the order of automorphism ϕ is a power of some prime number, then ϕ is an inner automorphism.*

Proof. According to Lemma 1.1, to prove the Theorem 2 it suffices to show that the equality $N^\phi = N$ holds for any normal subgroup $N \in \mathcal{M}_n$.

Suppose that there is a normal subgroup $A \in \mathcal{M}_n$ which is not ϕ-invariant, that is $A^\phi \neq A$. On the other hand, by Lemma 1.2 the centralizer of each such subgroup A is not trivial. Let p^r be the order of automorphism ϕ, where p is some prime number. The number p^r divides n by definition.

Since the subgroups of the cyclic group $\langle\phi\rangle$ of order p^r are linearly ordered by inclusion, one can choose some subgroup N with the minimal centralizer among all non-ϕ-invariant subgroups $A \in \mathcal{M}_n$. Being a subgroup of the group $\langle\phi\rangle$ of order p^r, this minimal nontrivial centralizer is generated by some automorphism of the form ϕ^{p^k}, where $1 < k < r$.

By virtue of the minimality, the subgroup $\langle\phi^{p^k}\rangle$ is contained in the centralizer of any subgroup $A \in \mathcal{M}_n$. Hence, the automorphism ϕ^{p^k} centralizes all subgroups $A \in \mathcal{M}_n$. According to

Lemma 1.1 we obtain that automorphism ϕ^{p^k} is inner.

We shall use the following lemma, that was proved in the paper [6].

LEMMA **2.1.** (see [6, Lemma 3]) *Let $\varphi : G \to G$ be an arbitrary automorphism and H be a normal subgroup of the group G such that the quotient group G/H is a non-abelian and simple. In that case, if the subgroups H, $H^{\varphi}, ..., H^{\varphi^{k-1}}$ are pairwise distinct and $H^{\varphi^k} = H$, then the quotient group $G/\bigcap\limits_{i=1}^{k} H^{\varphi^i}$ is decomposed into the direct product of normal subgroups $H_j/\bigcap\limits_{i=1}^{k} H^{\varphi^i}$, $j = 1, 2, ..., k$, wherein each quotient group $H_j/\bigcap\limits_{i=1}^{k} H^{\varphi^i}$ is isomorphic to G/H and $H_j = \bigcap\limits_{\substack{i=1 \\ i\neq j}}^{k} H^{\varphi^i}$.*

To use Lemma 2.1, suppose that $G = B(m, n)$, $\varphi = \phi$ and let $H = N$ be the normal subgroup of the group $B(m, n)$ with the minimal normalizer $\langle \phi^{p^k} \rangle$. The quotient group $B(m, n)/N$ is a Tarski monster, since $N \in \mathfrak{M}_n$. In particular, $B(m, n)/N$ is a non-abelian and simple group. By Lemma 2.1, the quotient group $B(m, n)/K$ is decomposed into the direct product of subgroups N_0/K, N_1/K, ... , N_{p^k-1}/K, where $K = \bigcap\limits_{i=0}^{p^k-1} N^{\phi^i}$.

As it was mentioned above, the automorphism ϕ^{p^k} is inner, that is for some element $u \in B(m, n)$ we have the equality $\phi^{p^k} = i_u$, where i_u is the inner automorphism generated by u. Since the automorphism ϕ^{p^k} has order p^{r-k}, the relation $(i_u)^{p^{r-k}}(x) = x$ holds for any $x \in B(m, n)$. In the other words the element $u^{p^{r-k}}$ belongs to the center of $B(m, n)$. From Adian's theorem on the triviality of the center of the group $B(m, n)$ (see [1, Theorem 3.4]) it follows the equality $u^{p^{r-k}} = 1$. Moreover, the element u has order p^{r-k} because the automorphism ϕ^{p^k} has order p^{r-k}.

For the element uK of the group $B(m, n)/K$ there exist uniquely defined elements u_0K, $u_1K, ..., u_{p^k-1}K$ from the subgroups N_0/K, N_1/K, ... , N_{p^k-1}/K respectively such that

$$uK = u_0K \cdot u_1K \cdots u_{p^k-1}K. \tag{1.1}$$

Note that the relation $u^{p^{r-k}} = 1$ implies immediately the relations $u_i^{p^{r-k}}K = K$ for all $i = 0, 1, ..., p^k - 1$. In particular, we have $u_0^{p^{r-k}}K = K$.

According to Lemma 2.1 the groups N_0/K and $B(m, n)/N$ are isomorphic. By definition of the set $\mathfrak{M}_n$ any element of the quotient group $B(m, n)/N$ is contained in some cyclic subgroup of order n (see [2, Proposition 5.2]). Therefore, there exists an element $a \in B(m, n)$ such that $aK \in N_0/K$ and the element au_0K has the order n.

Besides, the elements $u_1K, \cdots, u_{p^k-1}K$ commute with both of the elements aK and u_0K in $B(m, n)/K$ since $u_1K, \cdots, u_{p^k-1}K$ belong to the direct factors N_1/K, ... , N_{p^k-1}/K respectively. Hence, the equality (1.1) implies the relations

$$u^s a u^{-s} K = u_0^s a u_0^{-s} K \tag{1.2}$$

for all integers s. Since ϕ is a splitting automorphism, we have the equality $a\, a^{\phi} a^{\phi^2} \cdots a^{\phi^{n-1}} = 1$.

Taking into account ϕ-invariance of the subgroup K, we obtain that the equality

$$aK \cdot a^{\phi}K \cdot a^{\phi^2}K \cdots a^{\phi^{n-1}}K = K \tag{1.3}$$

holds in the quotient group $B(m,n)/K$.

Recall that according to choice of subgroup N we have $a^{\phi^{p^k}}K \in N_0/K$. This allows us to rewrite the equality (1.3) in the form

$$bK \cdot b^{\phi}K \cdot b^{\phi^2}K \cdots b^{\phi^{p^k-1}}K = K, \tag{1.4}$$

where

$$b = aa^{\phi^{p^k}}a^{\phi^{2p^k}} \cdots a^{\phi^{(n/p^k-1)p^k}}. \tag{1.5}$$

Moreover, it is easy to see that the elements bK, $b^{\phi}K$, $b^{\phi^2}K, \cdots$, $b^{\phi^{p^k-1}}K$ belong to the direct components N_0/K, N_1/K, ... , N_{p^k-1}/K respectively. Hence, the equality (1.4) immediately implies that all the factors on the left-hand side of the equality (1.4) are trivial. In particular, we have the equality $bK = K$.

Since $\phi^{p^k} = i_u$, one can rewrite the equality (1.5) in the form

$$b = a \cdot uau^{-1} \cdot u^2au^{-2} \cdot u^{n/p^k-1}au^{-(n/p^k-1)}.$$

Accordingly, we obtain equality

$$bK = aK \cdot uau^{-1}K \cdot u^2au^{-2}K \cdots u^{n/p^k-1}au^{-(n/p^k-1)}K$$

in the quotient group. Then using the relations (1.2), we get the equality

$$bK = a \cdot u_0au_0^{-1} \cdot u_0^2au_0^{-2} \cdots u_0^{n/p^k-1}au_0^{-(n/p^k-1)}K.$$

Next, we use the following identity

$$a \cdot u_0au_0^{-1} \cdot u_0^2au_0^{-2} \cdots u_0^{n/p^k-1}au_0^{-(n/p^k-1)} = (au_0)^{n/p^k} \cdot u_0^{-n/p^k}.$$

The number n/p^k is divided by p^{r-k} since n is divided by p^r. By virtue of the equality $u_0^{p^{r-k}}K = K$, we finally obtain $bK = (au_0)^{n/p^k}K$. Hence, we obtain $(au_0)^{n/p^k}K = K$ because $bK = K$. This contradicts to condition of choice of the the element a, according to which the element au_0K has order n in the group $B(m,n)/K$. Theorem is proved.

COROLLARY **2.1.** *If p is an odd prime number and $n = p^k \geq 1003$, then any splitting automorphism of period n of the group $B(m,n)$ is an inner automorphism.*

□

References

[1] S. I. Adian, *The Burnside problem and identities in groups*, Nauka, Moscow 1975; English transl., Ergeb. Math. Grenzgeb., vol. 95, Springer-Verlag, Berlin–New York 1979.

[2] S. I. Adyan; I. G. Lysenok, Groups, all of whose proper subgroups are finite cyclic. *Izv. Akad. Nauk SSSR Ser. Mat.* 55 (1991), no. 5, 933–990 (in russian); translation in Math. USSR-Izv. 39 (1992), no. 2, 905–957.

[3] V. S. Atabekyan, Simple and free periodic groups. *Vestnik Moskov. Univ. Ser. I Mat. Mekh.* (1987), no. 6, 76–78 (in russian); translation in Mosc. Univ. Math. Bull. 42(6) (1987) 80–82.

[4] V. S. Atabekyan, On periodic groups of odd period $n \geq 1003$. *Mat. Zametki*, 82:4 (2007), 495500 (in russian); translation in Math. Notes, 82:4 (2007) 443447.

[5] V. S. Atabekyan, Normal automorphisms of free Burnside groups. *Izv. RAN. Ser. Mat.* **75**:2 (2011) 3–18 (in russian); translation in Izv. Math. **75**:2 (2011) 223–237.

[6] V. S. Atabekyan, Splitting automorphisms of free Burnside groups. *Mat. Sb.*, **204**(2) (2013) 31–38 (in russian); translation in Sbornik: Mathematics, (2013), 204:2, 182–189.

[7] E. A. Cherepanov, Normal automorphisms of free Burnside groups of large odd exponents, *Internat. J. Algebra Comput* **16**(5) (2006) 839–847.

[8] O. H. Kegel, Die Nilpotenz der H_p-Gruppen, *Math. Z.*, v. 75 (1961), 373–376.

[9] E. I. Khukhro, Nilpotency of solvable groups admitting a splitting automorphism of prime order, *Algebra i Logika* 19:1 (1980), 118-129; English transl. in Algebra and Logic 19:1 (1980), 77-84.

[10] E. Jabara, Groups admitting a 4-splitting autmomorphism, *Rend. Circ. Mat. Palermo*, II. Ser., 45:1 (1996), 84–92.

[11] V. D. Mazurov, Yu. I. Merzlyakov and V. A. Chirkin (eds.), *The Kourovka notebook. Unsolved problems in group theory*, 11th ed., Institute of Mathematics, Siberian Branch of the USSR Academy of Sciences, Novosibirsk 1990. (Russian)

[12] A.Yu. Ol'shanskiĭ., Groups of bounded period with subgroups of prime order, *Algebra i Logika* 21:5 (1982), 553–618; English transl. in Algebra and Logic 21:5 (1982), 369–418.

[13] A. Yu. Olshanskii, *The Geometry of Defning Relations in Groups*. Kluwer Academic Publishers, 1991.